

Product List

A company that shares respect for life and beauty,
BioWellSpharm Co., Ltd., a company that gives health with love.



KGSP
Korean Good Supplying Practice
Certificate Holder





Biowellspharm OEM Products



A N E S T H E T I C S



엠마오 플러스크림

[국소마취크림]

Emmao Plus Cream
Local Anesthetics

Classification number	General 121	Insurance code	Non-payment	Packaging unit	450 g
Components /content	96mg lidocaine of 1g				
Efficacy /effectiveness	Decreased sensitivity of male genitals' tactile sensation				
Usage / capacity	Apply 0.24 to 0.65 g as lidocaine (23 to 62 mg as lidocaine) 5 to 15 minutes before sex. Capacity and application time vary according to individuals and use the lowest effective dose. Do not administer more than 2 g (184 mg as lidocaine) within 24 hours.				



엠마오 5%크림

[국소마취크림]

Emmao 5% Cream
Local Anesthetics

Classification number	Professional 121	Insurance code	Non-payment	Packaging unit	450 g
Components /content	25 mg of lidocaine and 25 mg of prilocaine in 1g				
Efficacy /effectiveness	1. Surface anesthesia of the skin during needle insertion or superficial surgical treatment 2. Reproductive mucosal surface anesthesia (e.g. before removing condylom)				
Usage / capacity	1. Skin 1) When inserting needles (e.g., intravenous intubation, blood collection), about 2 g shall be applied for at least 1 hour and up to 5 hours. 2) When performing superficial surgical procedures [e.g. extraction of molluscum] about 1.5~2 g/10cm ² shall be applied for at least one hour. 3) During superficial surgical major surgery (e.g., ruptured skin graft), 1.5~2g/10cm ² minimum of 2 hours and maximum of 5 hours shall be applied. 2. Genital mucosa When removing condylom, apply about 10 g to the treatment area for 5 to 15 minutes, remove the medicine, and immediately begin the procedure. * Infants aged 3 to 12 months: The total amount does not exceed 2 g for the 16cm ² of the applied part.				



리겐에스크림

[국소마취크림 / 수출용]

Regeness Cream
Local Anesthetics

Classification number	General 121	Insurance code	Non-payment	Packaging unit	450 g
Components /content	96mg lidocaine of 1g				
Efficacy /effectiveness	Decreased sensitivity of male genitals' tactile sensation				
Usage / capacity	Apply 0.24 to 0.65 g as lidocaine (23 to 62 mg as lidocaine) 5 to 15 minutes before sex. Capacity and application time vary according to individuals and use the lowest effective dose. Do not administer more than 2 g (184 mg as lidocaine) within 24 hours.				

VITAMIN B1



하프로킨주
[푸르셀티아민염산염]

Haprokin Inj.
Fursultiamine Hydrochloride

Classification number	Professional 312	Insurance code	Non-payment	Packaging unit	10 ml x 10Amps.												
Components /content	Fursultiamine Hydrochloride 5.46 mg/ml of 1 Amp (10 ml)																
Efficacy /effectiveness	<p>1. Prevention and treatment of vitamin B1 deficiency 2. Dissemination of vitamin B1 in insufficient intake from food due to increased demand (consumable diseases, thyroid gland dysfunction, pregnant women, lactating women, severe physical labor, etc.) 3. Wernicke encephalopathy 4. Cardiac beriberi disability 5. In the case that vitamin B1 deficiency or metabolic disorder caused by the following conditions is believed to be involved:</p> <table border="1" style="width: 100%; text-align: center;"> <tr> <td style="width: 16.6%;">1</td> <td style="width: 16.6%;">2</td> <td style="width: 16.6%;">3</td> <td style="width: 16.6%;">4</td> <td style="width: 16.6%;">5</td> <td style="width: 16.6%;">6</td> </tr> <tr> <td>neuralgia</td> <td>Muscle pain, joint pain.</td> <td>Peripheral neuropathy, peripheral nerve paralysis</td> <td>Myocardial metabolic disorder</td> <td>gastrointestinal such as constipation</td> <td>Postoperative paralyzing</td> </tr> </table> <p>It should not be used aimlessly for about a month without effect for deficiency of vitamin B1 or where metabolic disorders are presumed to be involved.</p>					1	2	3	4	5	6	neuralgia	Muscle pain, joint pain.	Peripheral neuropathy, peripheral nerve paralysis	Myocardial metabolic disorder	gastrointestinal such as constipation	Postoperative paralyzing
1	2	3	4	5	6												
neuralgia	Muscle pain, joint pain.	Peripheral neuropathy, peripheral nerve paralysis	Myocardial metabolic disorder	gastrointestinal such as constipation	Postoperative paralyzing												
Usage / capacity	Fursultiamine is usually administered in subcutaneous, intramuscular, or intravenous injections of 5 - 100 mg per adult per day. Increase and decrease according to symptoms																

VITAMIN D



본마린주
[콜레칼시페롤]

Bonmarin Inj.
Vitamin D3

Classification number	Professional 311	Insurance code	Non-payment	Packaging unit	1.5 ml x 10 Amps.
Components /content	Vitamin D3 7.5mg / 1.5 ml (5 mg/ml) (300,000 IU) of 1 Amp (1.5 ml)				
Efficacy /effectiveness	<p>1. Prevention and treatment of vitamin D deficiency in the elderly and adolescents who lack vitamin D 2. Treatment of vitamin D deficiency in adults with vitamin D deficiency</p>				
Usage / capacity	<p>Use it according to the doctor's prescription, but adjust the amount of 25-hydroxy vitamin D in the blood. Treatment effects and drug resistance should be considered within three months for adults and six months for adolescents after the drug is administered, and based on this, the decision is made to re-dose. Abnormal reactions increase with over-dosing, so do not administer more than necessary.</p> <ul style="list-style-type: none"> - Adult: Muscle injection of 200,000 IU as a Vitamin D3. - Adolescents: Muscle injection of 200,000 IU as a Vitamin D3. - Elderly: Muscle injection of 100,000 IU as a Vitamin D3. <p>The annual dose should not exceed 600,000 IU as a Vitamin D3.</p>				

ANTI-OXIDANT



글루치온주 600 mg
[글루타티온/환원형]

Gluthione Inj.
GLUTATHIONE (Reduced)

Classification number	Professional 399	Insurance code	Non-payment	Packaging unit	10 Vials
Components /content	GLUTATHIONE (Reduced) 600 mg				
Efficacy /effectiveness	Prevention of neurological diseases by cisplatin or similar-series chemotherapy				
Usage / capacity	Muscle injections or asymptotic intravenous intravenous injections of 600-1200 mg per day for severe cases. Muscle injection or asymptotic intravenous injection of 300 - 600 mg per day for treatment over several days or in mild cases.				



O T H E R S

엘디라제주 / 엘디엘주

[히알루로니다제]



Ldirase Inj. / LDL Inj.

Hyaluronidase 1500 IU



Actual size difference

Classification number	Professional 395	Insurance code	Non-payment	Packaging unit	10 Vials
Components /content	Hyaluronidase 1,500 IU of 1 Vial				
Efficacy /effectiveness	Increased penetration during subcutaneous or muscle injection, local anesthetic and subcutaneous injection Promote reabsorption of excess fluid and blood in tissues				
Usage / capacity	<p>Adults, children, and the elderly</p> <p>1. When subcutaneous injection (bulk subcutaneous injection) Dissolve 1,500 IU as a hyaluronidase in 1 ml of Sterile Water for Injection or physiologic saline solution and inject it into the affected area before subcutaneous injection, or into a tube 2 cm above the infusion needle at the beginning of injection. This approximately 1,500 IU is appropriate for administration of 500 to 1,000 ml of sap. Children and the elderly should carefully control the speed and total dose when administering the sap, especially if they have a renal disorder, to avoid excess moisture.</p> <p>2. When hypodermic or muscle injections It is used by directly dissolving about 1,500 IU into the injection liquid to be administered.</p> <p>3. Local anesthetic It is used by dissolving it in a local anesthetic injection to administer about 1,500 IU. When used in ophthalmology, a concentration of 15 IU per ml is recommended.</p> <p>4. Extravascular effusion In cases of diffusion rather than local cases, after the appearance of extravascular effusion, about 1,500 IU is dissolved in 1 ml of the main use water or physiological saline solution as soon as possible and infiltrates the lesion area.</p> <p>5. hematoma Dissolve about 1,500 IU in the Sterile Water for Injection or 1 ml of the saline liquid to the affected area. The drug is dissolved in an injection that will be administered with approximately 1 ml of main water immediately prior to use. Subcutaneous injection should appear with extracellular fluid in the body. The drug can be mixed with commonly used transfusion agents. Examples of use in mass hypodermic injections with physiological saline solution, 4 percent glucose of 0.18 percent sodium chloride, 2.5 percent glucose of 0.45 percent sodium chloride, and 5 percent glucose injection were reported. Calium is administered in 34 mmol/L to the injection of glucose or physiologic saline. A sap containing electrolytes is more suitable than a sap containing no electrolytes and should not be injected too quickly. The drug is used in combination with morphine, diamorphine, hydromorphine, chlorpromazine, metoclopramide, promazine, dexamethasone, local anesthetic and epinephrine.</p>				



엘디엘정


















[알긴산, 카르복시메틸셀룰로오스나트륨]

















LDL Tab.





Alginate acid & Carboxymethylcellulose Sodium

Classification number	Professional 399	Insurance code	Non-payment	Packaging unit	300 T.
Components /content	Alginate acid 200 mg, Carboxymethylcellulose Sodium 100 mg of 1 tab				
Efficacy /effectiveness	A supplement to weight loss by reducing food intake				
Usage / capacity	You can take two tablets three times a day before meals, and if necessary, you can take two tablets before bedtime. It is recommended to take it with sufficient water. It is recommended to take the recommended dose for at least three to four weeks.				

Product	Name	Unit	Main Components	Efficacy and Effectiveness	Note
VITAMIN					
	Haprokin Inj	10A	Fursultiamine Hydrochloride 5.46 mg/ml of 1 Amp (10 ml)	Prevention and treatment of vitamin B1 deficiency	Vitamin B1 (Garlic Injection)
	Fursultiamine Inj	10A/50A			
	Bonmarin Inj.	10A	Vitamin D3 7.5 mg (300,000 IU) of 1 ampule (1.5 ml)	Prevention and treatment of vitamin D deficiency in the elderly and adolescents who lack vitamin D	
	Vita-D Inj	5A	Vitamin D3 5 mg (200,000 IU) of 1 ampule (1 ml)	Prevention and treatment of vitamin D deficiency in the elderly and adolescents who lack vitamin D	Vitamin D3
	BMHeidi Inj.	10V			
	Vitavella Inj	10PF			
	MultiVita Inj.	10V	Among 1 vials Vitamin A Vitamin B. Vitamin C Vitamin D Vitamin E	1.Vitamin maintenance therapy for patients who are nourished with sap 2.Vitamins for the following patients (excluding vitamin K) 3.Deficiency Prevention: Severe burns, fractures, other trauma, severe infectious disease coma during surgery	Multi Vitamin I njection.
	EsmultiVita Inj.				
	VitaminC Inj.	20 mL x 10V	Ascorbic acid 10 g of 1 vial (20 ml)	Deficiency and prevention and treatment of vitamin C	High capacity vitamin C
	Ascovic Inj.				
	BCup Inj.				
	Livovita Inj.	1 ml x 50A	Riboflavin. 12.71 mg/ml	Prevention and treatment of vitamin B2 deficiency	Vitamin B2
	Hiv5 Inj.	50A	D-Panthenol 250 mg/ml	Prevention and treatment of pantothenic acid deficiency	Vitamin B5
	Vitamo Inj.				
	Hiv6 Inj.	50A	Pyridoxine hydrochloride 50 mg/ml	Prevention and Treatment of Vitamin B6 Deficiency	Vitamin B6
	Piridoxin Inj				
	Panbicomp Inj	4 ml x 10A	Thiamine hydrochloride, riboflavin, nicotinic acid, dexpantenol, pyridoxin, biotin, cyanocobalamin	Supplied B vitamins.	Vitamin B-complex

Product	Name	Unit	Main Components	Efficacy and Effectiveness	Note
VITAMIN					
	HaikoMin Inj.	2 ml x 50A	Hydroxocobalamin 5 mg of 1 ampule (2 ml)	Acute neuropathy, multiple neuropathy, sciatica, rheumatoid neuropathy, sclerosis, scapularis, cancer and trauma-induced neuralgia, shingles, and migraine -induced neuropathy	Vitamin B12
	Red-up Inj.				
	Hikoval Inj.				
PLACENTA					
	Linec Inj.	2 ml x 50A	Hominis Placenta hydrolysate 2 ml of 1 ampule (2 ml)	Improvement of liver function in chronic liver disease	Placenta injection
	Mersmon Inj.	2 ml x 50A	Hominis Placenta extract 2 ml of 1 ampule (2 ml)	Improvement of Menopausal Disorder	
	Melsmon Inj.				
	Lizube Inj.	50A			
ANTIOXIDANT					
	Gluthione Inj. 600mg	10V	GLUTATHIONE (Reduced) 600 mg of 1 Vial	Prevention of neurological diseases by cisplatin or similar-series chemotherapy	Whitening injection
	Detoxion Inj. 1200mg	10V	GLUTATHIONE (Reduced) 1200 mg of 1 Vial		
	Thioctacid Inj.	24 ml x 5A	Tromethamine 39.7 mg/ml (Insurance Code - 642200991 4,866/A)	Alleviation of diabetic multiple neuropathy	
	Idela Inj.	10A	Thioctic Acid 25 mg of 1 ampule (5ml)	Dissemination of cases where demand for thioctic acid increases (in severe physical labor) Irrational hearing loss due to Leigh syndrome (subacute necrotic cerebrospinal inflammation) poisoning (streptomycin, kanamycin) and noise tolerance (occupational)	Cinderella Injection
	Cindella Inj.				
LOCAL ANESTHETICS					
	Emmao Plus Cream	450 g	96mg lidocaine of 1g	Decreased sensitivity of male genitals' tactile sensation 25 mg of lidocaine and 25 mg of prilocaine in 1g	Anesthesia cream
	Lidokin cream	30 g			
	Dermacan Cream	500 g			
	Emmao 5% Cream	450 g	25 mg of lidocaine and 25 mg of prilocaine in 1g	1. Surface anesthesia of the skin during needle insertion or superficial surgical treatment 2. Reproductive mucosal surface anesthesia (e.g. before removing condylom)	
	Regeness Cream (For export)	500 g	105.6 mg of 1g lidocaine	Decreased sensitivity of male genitals' tactile sensation	

Product	Name	Unit	Main Components	Efficacy & Effectiveness	Note
MINERAL					
	Multiblue 5 Inj.	10 ml x 25V	Selenium 200 µg, Zinc 10 mg, copper 4 mg, Manganese 1 mg, chromium 40 µg of 1 Vial (10 ml)	Mild elements in the distribution of meridians Dissemination of (zinc, copper, manganese, selenium, chromium)	Essential minerals
	Jaeil Mineral 5 Inj.	10 ml x 10V			
	Masi Inj. 10%	5 ml x 25A	Magnesium sulfate hydrate 0.1 g/ml	Cramps, seizures of narcolepsy, cervical cyst prevention, electrolyte dissemination.	Magnesium Injection (Mg)
	Selenu Inj.	10 ml x 10V	Selenium 400 µg of 1 Vial (10 ml)	Replenishment of selenium during parenteral alimentation (TPN)	Selenium Injection (Se)
	Seleniumin Inj.				
	Zinkin Inj.	10 ml x 10V 20 ml x 10V	Zinc 1 mg/ml of 1 vial	Replenishment of Zinc during parenteral alimentation (TPN)	Zinc (Zn) Injection
	Zinkon Inj.	10 ml x 10V			
AMINO ACID					
	Nutrihex Inj	100 ml x 20EA 250 ml x 10EA	L-tryptophane, L-histidine, L-isoryushin, arginine, glycine, alanine, prolene (body L), L-metionine, L-treonine, L-phenylalanine L-valin, L-Rucine, etc.	Amino Acid Water Solution (Verbital Injection Foundation Nutrient)	Amino acid solution
	Winnapferry Inj	362 ml	7.3 g of L-Lucine, 7.25 g of L-Lucine Hydrochloride and more	Dissemination of moisture, electrolytes, amino acids, calories, essential fatty acids and omega-3 fatty acids to patients who are unable, insufficiently or prohibited from taking nutrition in the oral or gastrointestinal tract.	3-chamber
	Vitasol Inj	250 ml x 10ea	L-isomericin, L-Lucine, L-Lucin, xylitol, sodium riboflavinphosphate, pyridoxin hydrochloride, nicotinic acid mide, inositol and others	Amino Acid Water Solution (Verbital Injection Foundation Nutrient)	Amino acid solution
	Daehan El Arginine	100 ml x 10 Bottle 150 ml x 10 Bottle	L-Arginine Hydrochloride 100 mg/ml of 1 Bottle	Birth factor circuit abnormality, lysine uroprotein intolerance	Arginine injection
	Dipeptiben Inj.	50 ml x 10 Bot.	N(2) - Elalanyl-Elglutamine	Glutamine distribution	Glutamine
PDRN					
	Placentex Inj.	3 ml x 5A	5.625 mg of sodium polydeoxyribonucleotide among 1 ampoule (3 ml)	Treatment of wounds and tissue recovery due to skin transplantation.	Salmon injection (PDRN)
	Heidial Inj.	3 ml x 10V			
GENERAL ANESTHETICS					
	1% of MyeongMun provive Inj.	10 mlx10V 20 mlx5V 50 mlx1V	Propofol 10 mg/ml (Insurance code : 10 ml 649805211 1,317/V) (Insurance code: 20 ml 649805221 4,061/V) (Insurance code: 50 ml 649805231 9,619/V)	Induction and maintenance of general anesthesia for adults and children aged 3 years or older; Sedation of a heavily ventilated patient. Sedation of consciousness during surgery.	A general anesthetic
	DongKuk Popol Inj.	8 mlx10A 12 mlx5A 20 mlx5A			

Product	Name	Unit	Main Components	Efficacy & Effectiveness	Note
GROWTH HORMONE					
	Caretropine Inj.	1.5 ml x 4EA	Somatropin 75 mg (22.5 IU) of 1 Vial(1.5ml)	Alternative treatment for growth hormone in adults with growth hormone deficiency identified by two dynamic tests	Adult growth hormone
OTHERS					
	Ldirase Inj.	10V	Hyaluronidase 1,500 IU of 1 Vial	Increased penetration during subcutaneous injections, muscle injections, local anesthetics and subcutaneous injections and facilitate reabsorption of excess fluid and blood in tissues	Pain (corrosion) management Filler Removal
	LDL Inj.	10V			
	Liporaje Inj.	10V			
	Petlise Inj.	25A	L-Carnitine 1 g of 1 ampule (5ml)	Primary, secondary carnitin deficiency, angina, acute myocardial infarction, terminal stage of hemodialysis, the treatment of carnitin deficiency of renal disease patient	Elkarnitine injection
	Kartin Inj.	10A			
	Atipi Inj.	2 ml x 50A	Adenosine Triphosphate dinatrium trihydrate 21.961 mg of 1 ampule (2ml)	Improving liver function in work force, heart failure, chronic hepatitis, aftereffects of head trauma, and controllable stable fatigue.	Nucleic acid injection
	Glucoline Es Inj.	20 ml x 50A	Elcysteine hydrochloride 20 mg, Monoammonium glycirizinate 53 mg, Aminochoic cid 400 mg of 1 ampule (20ml)	Improvement of urticaria, eczema, allergic skin disease, adjuvant therapy of drug addiction, and liver function of chronic diseases	Licorice injection
	Same Inj.	5V	S-Adenosyl-L-Methionine sulfate toxylate 192 mg	Depression	S-Adenyl-l-Methionine
	Imualpha Inj.	2V	Cymosin Alpha 1 1.6 mg	Supplementary therapy for influenza vaccination in older patients with impaired immune function	Immunization injection
	Hiront Inj.	10Tube	Sodium hyaluronic acid 10 mg/ml	Deformed systolic arthritis, scapular gastritis	A water-glossi injection
	Escolin Inj.	10A	Colin Alpocerate 250 mg/ml	Secondary symptoms and degeneration or degenerative cerebral matrix psychosis caused by cerebrovascular deficits	Improve brain function
	Mezzopilin Inj.	10A	Pentoxifiline 20 mg	Cerebral circulation disorder, blood flow circulation in the eyes, peripheral arterial circulation disorder	Peripheral vascular dilatation brain functional enhancer
	Ginko Inj.	5 ml x 25A	Ginkgo Biloba 17.5 mg of 1 ampule (5 ml)	Decrease in intellectual and alert ability, dizziness, tinnitus, headache, vision impairment, memory loss, anxiety, depression, hearing and language skills due to dyslexia, vascularity, cervical symptoms, etc.	Ginkgo Biloba X
	Nepopen Inj.	20A	Nefopam 20 mg 1 ampule (2 ml)	Acute pain hypertrophy, especially postoperative pain	Pain injection
	Eldiel Inj.	300T	alginic acid 200 mg, carboxymethyl cellulose 100 mg of 1 tablet	A supplement to weight loss by reducing food intake	Weight Loss Supplement